

Lanthanide-doped CaF_2 and SrF_2 nanoparticles for biomedical applications: in vivo and in vitro experimental studies

Corinne Portioli¹, Marco Pedroni², Donatella Benati¹, Stefano Dusi³, Marta Donini³, Raffaella Mariotti¹, Roberta Bonafede¹, Luigi Perbellini⁴, Marzia Cerpelloni⁴, Adolfo Speghini², Marina Bentivoglio¹

¹ Dipartimento di Scienze Neurologiche e del Movimento, Università di Verona, Verona, Italy - ² Dipartimento di Biotecnologie, Università di Verona, Verona, Italy - ³ Dipartimento di Patologia, Università di Verona, Verona, Italy - ⁴ Dipartimento di Salute Pubblica e di Comunità, Università di Verona, Verona, Italy

Among the wide range of nanoparticles (NPs) studied for diagnostic and therapeutic applications lanthanide-doped nanosystems have raised special interest [1]. Their very small dimension (10 nm) and upconversion emission property have increased the range of their applications from contrast agent probes in bioimaging to drug delivery systems [2,3]. Here, the cytotoxicity of rare earth (Yb and Er)-doped CaF_2 and SrF_2 NPs has been investigated both *in vitro* and *in vivo*. *In vitro* studies have been conducted in a motoneuron cell line as model of neuronal interaction, and in a line of human dendritic cells which play a key role in the immune response. In the motoneuron cell line, a weak response was observed at early time points while the cell viability showed an increment, except for the highest concentration of lanthanide-doped NPs. The levels of cytokines released from human dendritic cells were low and dose-dependent. The NP biodistribution was investigated after a single peripheral administration in mice. Aggregates of NPs were shown, with different techniques, mostly in peripheral organs (spleen and liver) after one day. A limited penetration of both CaF_2 and SrF_2 NPs was seen in the brain parenchyma, associated with a mild astrocytic activation. Since the present *in vitro* findings indicate that lanthanide-doped NPs are safe, and the *in vivo* data show that they can enter the brain parenchyma crossing the blood-brain barrier, these NPs may represent promising tools for diagnostic and therapeutical applications.

Supported by The Fondazione Cariverona ("Verona Nanomedicine Initiative").

References

- [1] Chatterjee et al. (2008) Upconversion fluorescence imaging of cells and small animals using lanthanide doped nanocrystals. *Biomaterials* 29; doi:10.1016/j.biomaterials.2007.10.051.
- [2] Bae et al., (2012) Endocytosis, intracellular transport, and exocytosis of lanthanide-doped upconverting nanoparticles in single living cells. *Biomaterials* 33, doi:10.1016/j.biomaterials.2012.08.039.
- [3] Ni et al., (2014) Dual-targeting upconversion nanoprobe across the blood-brain barrier for magnetic resonance/fluorescence imaging of intracranial glioblastoma. *ACS Nano* 8; doi: 10.1021/nl406197c.

Keywords

Lanthanide ions; Nanoparticles; Cell viability; Biodistribution.